TUM-T31-92/95 hep-ph/9510383

arXiv:hep-ph/9510383v1 24 Oct 1995

PHENOMENOLOGY OF ε_K BEYOND LEADING LOGARITHMS*

Ulrich Nierste†

Physik-Department, Techn. Univ. München, D-85747 Garching, Germany

I present the QCD short distance coefficient η_3 of the $\Delta S = 2$ hamiltonian in the *next-to-leading order* (NLO) of renormalization group improved perturbation theory. Since now all QCD-factors η_1, η_2 and η_3 are known with NLO accuracy, a much higher precision in the analysis of ε_K can be achieved. The CKM phase δ, $|V_{td}|$ and the mass difference Δm_{B_s} in the $\mathrm{B}_s^0 - \overline{\mathrm{B}}_s^0$-system are predicted from the measured values of ε_K and Δm_{B_d}. Finally I briefly look at the $\mathrm{K_L} - \mathrm{K_S}$-mass difference. This work has been done in collaboration with Stefan Herrlich.

1. The $|\Delta S| = 2$-hamiltonian in the NLO

The low-energy hamiltonian inducing $\mathrm{K}^0 - \overline{\mathrm{K}^0}$-mixing reads:

$$H^{|\Delta S|=2} = \frac{G_F^2}{16\pi^2} M_W^2 \left[\lambda_c^2 \eta_1^\star \frac{m_c^{\star\,2}}{M_W^2} + \lambda_t^2 \eta_2^\star S(\frac{m_t^{\star\,2}}{M_W^2}) + 2\lambda_c\lambda_t \eta_3^\star S(\frac{m_c^{\star\,2}}{M_W^2}, \frac{m_t^{\star\,2}}{M_W^2}) \right] b(\mu) Q_{S2}(\mu) + \text{h.c.} \qquad (1)$$

Here G_F is the Fermi constant, M_W is the W boson mass, $\lambda_j = V_{jd} V_{js}^*$ comprises the CKM-factors and Q_{S2} is the local $|\Delta S| = 2$ four-quark operator

$$Q_{S2} = (\overline{s}_j \gamma_\mu (1-\gamma_5) d_j)(\overline{s}_k \gamma^\mu (1-\gamma_5) d_k) \qquad (2)$$

with j and k being colour indices. $m_q^\star = m_q(m_q)$, $q = c, t$, are running quark masses in the $\overline{\mathrm{MS}}$ scheme. The Inami-Lim functions $S(x), S(x,y)$ describe the $|\Delta S| = 2$-transition amplitude in the absence of strong interaction. The short distance QCD corrections are comprised in the coefficients η_1, η_2 and η_3 with a common factor $b(\mu)$ split off. They are functions of the charm and top quark masses and of the QCD scale parameter Λ_{QCD}. Further they depend on the *definition* of the quark masses used in the Inami-Lim functions: In (1) the η_i's are defined with respect to $\overline{\mathrm{MS}}$ masses $m_q^\star$ and are therefore marked with a star.

With actual values of the input data the results of the old *leading log approximation* read

$$\eta_1^{\star\,\mathrm{LO}} \approx 0.80\,, \qquad \eta_2^{\star\,\mathrm{LO}} \approx 0.62\,, \qquad \eta_3^{\star\,\mathrm{LO}} \approx 0.36\,. \qquad (3)$$

Now the NLO values read:

$$\eta_1^\star = 1.32^{\,+0.21}_{\,-0.23}\,, \quad \eta_2^\star = 0.57^{\,+0.00}_{\,-0.01}\,, \quad \eta_3^\star = 0.47^{\,+0.03}_{\,-0.04}\,, \qquad (4)$$

where $m_c^\star = 1.3\,\mathrm{GeV}$ and $\Lambda_{\overline{\mathrm{MS}}}^{\mathrm{NLO}} = 0.310\,\mathrm{GeV}$ has been used. η_2 and η_3 are almost independent of the input parameters.

The NLO calculation of η_2 has been performed in [1] and the NLO result for η_1 can be found in [2]. η_3 in (4) is new. Details of the calculation are presented in [3]. A phenomenological analysis using the NLO η_i's has been published in [4].

(3) and (4) clearly show the sizeable numerical effect of the NLO correction. Further there are conceptual reasons for going to the NLO:

i) The fundamental QCD scale parameter $\Lambda_{\overline{\mathrm{MS}}}$ cannot be used in LO calculations.

ii) The quark mass dependence of the η_i's is not accurately reproduced by the LO expressions. Especially the m_t-dependent terms in $\eta_3 \cdot S(m_c^{\star\,2}/M_W^2, m_t^{\star\,2}/M_W^2)$ belong to the NLO.

iii) Likewise the proper definition of the quark masses is a NLO issue: One must go to the NLO to learn how to use m_t measured at FERMILAB in low energy expressions such as (1). In NLO the $\overline{\mathrm{MS}}$ mass $m_t^\star$ is smaller than the *pole mass* m_t^{pole} by 8 GeV. It has been discussed at this conference to which definition of m_t the quoted CDF and D0 results for m_t refer. Presumably the measured quantity is m_t^{pole}.

iv) In the NLO the large LO error bars caused by renormalization scale dependences are reduced.

2. Phenomenology of ε_K

Let us first recall our present knowledge about the CKM matrix V_{CKM}: The precise measurements of $|V_{ud}|$ and $|V_{us}|$ also constrain $|V_{cd}|$, $|V_{cs}|$ and $|V_{tb}|$ via the unitarity

*Talk presented at the EPS-HEP 95 conference, Brussels, Jul. 27-Aug. 2 1995.

†supported by BMBW under contract no. 06-TM-743

of V_{CKM}. $|V_{cb}|$ is expected to lie in the range

$$0.037 \leq |V_{cb}| \leq 0.043 \tag{5}$$

After fixing $|V_{cb}|$ unitarity likewise pins down $|V_{ts}|$. We will further need

$$0.06 \leq \frac{|V_{ub}|}{|V_{cb}|} \leq 0.10. \tag{6}$$

Yet even for fixed $|V_{cb}|$ and $|V_{ub}/V_{cb}|$ the magnitude of the remaining CKM element $|V_{td}|$ is a sensitive function of the phase δ. Hence Δm_{B_d} and $|\varepsilon_K|$ provide complementary information on $|V_{td}|$: The former determines this element directly and the latter indirectly through δ.
Apart from $|V_{cb}|$ and $|V_{ub}/V_{cb}|$ two other key parameters are involved: The actual value $m_t^{\rm pole} = (180 \pm 12)\,\mathrm{GeV}$ for the top quark pole mass corresponds to

$$160\,\mathrm{GeV} \leq m_t^\star \leq 184\,\mathrm{GeV}. \tag{7}$$

Finally the hadronic matrix element of Q_{S2} in (2) is parametrized as

$$\langle \overline{\mathrm{K}^0}|Q_{S2}(\mu)|\mathrm{K}^0\rangle \;=\; \frac{8}{3} f_K^2 m_K^2 B_K/b(\mu).$$

We take

$$0.65 \leq B_K \leq 0.85, \tag{8}$$

which reflects the $1/N_c$ result as well as the ballpark of the lattice determinations.
The uncertainties in the input parameters other than $|V_{cb}|$, $|V_{ub}|/|V_{cb}|$, m_t and B_K do not significantly affect the analysis (see [4]). Using the LO η_i's, however, imposes an error onto ε_K, which is roughly of the same order as the one due to (5-8). The NLO shift in η_3 affects ε_K as much as pushing B_K from 0.75 to 0.82. This uncertainty has been neglected in most phenomenological analyses.
After fixing three of the key parameters the measured value for ε_K yields a lower bound at the fourth one. This feature has been used in the pre-top era to constrain m_t. Yet today one should focus on the CKM elements instead. The allowed range for $(|V_{cb}|, |V_{ub}/V_{cb}|)$ is shown in fig. 1.

Next δ is obtained from a simultaneous analysis of ε_K and Δm_{B_d}: For $|V_{ub}|/|V_{cb}| = 0.08, |V_{cb}| = 0.04, m_t^\star = 172\,\mathrm{GeV}, B_K = 0.75$ one finds the two solutions:

$$\delta^{\rm low} = 85^\circ, \qquad \delta^{\rm high} = 121^\circ. \tag{9}$$

They correspond to

$$|V_{td}|^{\rm low} = 9.1 \cdot 10^{-3}, \qquad |V_{td}|^{\rm high} = 10.7 \cdot 10^{-3}. \tag{10}$$

Accounting for the errors (5-8) leads to

$$47^\circ \leq \delta \leq 135^\circ, \qquad 7.3 \leq |V_{td}| \cdot 10^3 \leq 11.9, \tag{11}$$



Figure 1: For each pair $(m_t^\star, B_K)$ the measured value for ε_K defines a curve. The points below the curve are excluded. The rectangle limits the allowed range for $|V_{cb}|$ and $|V_{ub}/V_{cb}|$ obtained from tree-level b-decays according to (5) and (6).

where the upper bounds stem from $\Delta m_{B_d} = (0.470 \pm 0.0025)\,ps^{-1}$. Here $F_{B_d}\sqrt{B_{B_d}} = (195 \pm 45)\,\mathrm{MeV}$ has been used.
The ratio $\Delta m_{B_d}/\Delta m_{B_s}$ is theoretically much better understood than the mass differences separately. We can use the result (11) for $|V_{td}|$ to predict Δm_{B_s}:

$$6.2\,ps^{-1} \leq \Delta m_{B_s} \leq 26\,ps^{-1}. \tag{12}$$

More details can be found in [4], where slightly different values for m_t and Δm_{B_d} have been used.

3. The $\mathbf{K_L - K_S}$-mass difference

In the 1980s it was generally believed that long distance interactions make up more than half of the observed $\mathrm{K_L - K_S}$-mass difference. Today the picture has changed due to the NLO enhancement of η_1 and η_3 and the larger value for $\Lambda_{\overline{\rm MS}}$, which pushes η_1 further up. One finds

$$\frac{(\Delta m_K)_{\rm SD}}{(\Delta m_K)_{\rm exp}} \;=\; 0.7 \pm 0.2 \tag{13}$$

exhibiting a short distance dominance. This is in agreement with naive power counting: One expects the long distance part to be suppressed with a factor of $(\Lambda_{had}/m_c)^2$ compared to (13). Here Λ_{had} is a hadronic scale.

References

[1] A. J. Buras, M. Jamin and P. H. Weisz, Nucl. Phys. B347 (1990) 491.

[2] S. Herrlich and U. Nierste, Nucl. Phys. B419 (1994) 292.

[3] S. Herrlich and U. Nierste, *The complete* $|\Delta S| = 2$*–hamiltonian in the next-to-leading order*, preprint TUM-T31-86/95 in progress.
U. Nierste, *Indirect CP-Violation in the Neutral Kaon System Beyond Leading Logarithms and Related Topics*, thesis TU München, 1995, hep-ph/9510323.

[4] S. Herrlich and U. Nierste, preprint TUM-T31-81/95, hep-ph/9507262, Phys. Rev. D, in press.